

The rare K-decays and Z^0 -penguin contributions in the Topcolor-assisted technicolor models

Zhenjun Xiao^(1,2,3)*, Chongsheng Li^(1,2), Kuangta Chao^(1,2)

1. CCAST(World Laboratory) P.O. Box 8730, Beijing 100080, P.R.China
2. Department of Physics, Peking University, Beijing, 100871 P.R. China.
3. Department of Physics, Henan Normal University, Xinxiang, 453002 P.R. China.

February 12, 1999

Abstract

We calculate the new contributions to the rare decays $K^+ \rightarrow \pi^+ \nu \bar{\nu}$, $K_L \rightarrow \pi^0 \nu \bar{\nu}$ and $K_L \rightarrow \mu^+ \mu^-$ from new Z^0 -penguin and box diagrams induced by the unit-charged scalars ($\tilde{\pi}^\pm, \tilde{H}^\pm, \pi_1^\pm, \pi_8^\pm$) appeared in the Topcolor-assisted technicolor (TC2) models. We find that: (a) the unit-charged top-pion $\tilde{\pi}^\pm$ and b-pion $\tilde{H}^\pm$ can provide large contributions to the rare K-decays if they are relatively light; (b) the size of mixing elements $D_{L,R}^{ij}$ ($i \neq j$) is strongly constrained by the data of B^0 meson mixing: $|a_R^{ts}|, |a_R^{td}| < 0.01$ for $a_L^{td} = a_L^{ts} = 1/2$ and $m_{\tilde{H}^0} \leq 600 GeV$; (c) the enhancements to the branching ratios of rare K-decays from new scalars can be as large as one order of magnitude; (d) there is a strong cancellation between the short- and long-distance dispersive part of the decay $K_L \rightarrow \mu^+ \mu^-$, the constraint on the new short-distance part from this decay mode is thus not strong; (e) the typical TC2 model under study is generally consistent with the available rare K-decay data.

PACS numbers: 12.60.Nz, 12.15.Ji, 13.20.Jf, 13.40.Hq

*E-mail: zxiao@ibm320h.phy.pku.edu.cn; and dphnu@public.zz.ha.cn

1 Introduction

As is well-known, the study of loop effects can open an important window on electroweak symmetry breaking and physics beyond the standard model(SM). The examination of indirect effects of new physics in flavor changing neutral current (FCNC) processes in rare K- and B-decays [1, 2, 3, 4, 5] offers a complementary approach to the search for direct production of new particles at high energy colliders.

In the SM, the rare K-decays $K^+ \rightarrow \pi^+ \nu \bar{\nu}$, $K_L \rightarrow \pi^0 \nu \bar{\nu}$ and $K_L \rightarrow \mu^+ \mu^-$ are all loop-induced semileptonic FCNC processes determined by Z^0 -penguin and W-box diagrams[1]. Since these rare decay modes are theoretically clean and highly suppressed in the SM, they may serve as the good hunting ground for the new physics beyond the SM. Furthermore, the relevant experimental measurements are now reach to a reasonable or even high sensitivity[6, 7], which will help us to test or constrain the new physics models through studies about these rare decay modes.

In refs.[2, 3], the authors studied the FCNC effects on the mixing and rare decays for the K and B meson systems in the minimal supersymmetric standard model and the two Higgs doublet model. In this paper we will investigate the new contributions to the rare K-decays from the Z^0 -penguin diagrams induced by the unit-charged scalars appeared in the TC2 model [8].

Technicolor (TC) [9, 10] is one of the important candidates for the mechanism of naturally breaking electroweak symmetry. To generate ordinary fermion masses, extended technicolor (ETC) [11] models have been proposed. In walking technicolor theories[12], the large FCNC problem can be resolved and the fermion masses can be increased significantly[12]. The S parameter can be small or even negative in the walking technicolor models[13]. To explain the large hierarchy of the quark masses, multiscale walking technicolor models (MWTCM) are further constructed [14]. In order to generate large top quark mass without running afoul of either experimental constraints from the ρ parameter and the R_b data, the TC2 models was constructed recently [8, 15, 16].

In TC2 model, the relatively light top-pions ($\tilde{\pi}^\pm, \tilde{\pi}^0$), b-pions ($\tilde{H}^\pm, \tilde{H}^0, \tilde{A}^0$) and other bound states, may provide potentially large loop effects in low energy observables. This is the main motivation for us to investigate the contributions to the rare K-decays from the penguin and box diagrams induced by the internal exchanges of unit-charged top-pions, b-pions and technipions.

In a previous paper [4], we calculated the Z^0 -penguin contributions to the rare K-decays from technipions $\pi_1^\pm$ and $\pi_8^\pm$ in the MWTCM[14] and found that this model was strongly disfavored by the relevant data.

In this paper we calculate the new Z^0 -penguin contributions to the rare K-decays from the top-pions $\tilde{\pi}^\pm$, b-pions $\tilde{H}^\pm$ and technipions $\pi_1^\pm$ and $\pi_8^\pm$ appeared in the TC2 model [8]. We firstly evaluate the Z^0 -penguin and box diagrams induced by the unit-charged scalars, compare the relevant analytical expressions of effective couplings with the corresponding expressions in the SM, separate the new functions $C_0(\pi_i)$ and $C_{NL}\pi_i$ ($\pi_i = \tilde{\pi}^\pm, \tilde{H}^\pm, \pi_1^\pm, \pi_8^\pm$) which describe the effects of the new particles, and finally combine the new functions with their counterparts in the SM and use them directly in the calculation for specific decay modes.

This paper is organized as follows. In Sec.2 we briefly review the basic structures of the TC2 models and study the experimental constraints on the mixing matrices D_L and D_R . In Sec.3 we firstly show the standard model predictions for the branching ratios of rare K-decays, and then evaluate the new one-loop Feynman diagrams and extract out the new effective Z^0 -penguin couplings induced by the exchanges of unit-charged scalars. In following two sections, we present the numerical results for the branching ratios $B(K^+ \rightarrow \pi^+ \nu \bar{\nu})$, $B(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and $B(K_L \rightarrow \mu^+ \mu^-)_{SD}$ with the inclusion of new physics effects, and compare the theoretical predictions with the available data. The conclusions and discussions are included in the final section.

2 Basic structure of TC2 models

Apart from some differences in group structure and/or particle contents, all TC2 models [8, 15, 16] have the following common features: (a) strong Topcolor interactions, broken near 1 TeV, induce a large top condensate and all but a few GeV of the top quark mass, but contribute little to electroweak symmetry breaking; (b) TC interactions are responsible for electroweak symmetry breaking, and ETC interactions generate the hard masses of all quarks and leptons, except that of the top quarks; (c) there exist top-pions $\tilde{\pi}^\pm$ and $\tilde{\pi}^0$ with a decay constant $f_{\tilde{\pi}} \approx 50$ GeV. In this paper we will chose the most frequently studied TC2-I model[8] ¹ as the typical TC2 model to estimate the contributions to the rare K-decays in question from the relatively light unit-charged

¹In this paper, we use the term “TC2-I” model to denote the TC2 model constructed by Hill[8].

scalars. It is straightforward to extend the studies in this paper to other TC2 models.

2.1 TC2-I model, couplings and mass spectrum

In the TC2-I model[8] the dynamics at the scale ~ 1 TeV involves the following structure:

$$SU(3)_1 \times U(1)_{Y1} \times SU(3)_2 \times U(1)_{Y2} \times SU(2)_L \rightarrow SU(3)_{QCD} \times U(1)_{EM} \quad (1)$$

where $SU(3)_1 \times U(1)_{Y1}$ ($SU(3)_2 \times U(1)_{Y2}$) generally couples preferentially to the third (first and second) generation fermions. The breaking (1) typically leaves a residual global symmetry, $SU(3)' \times U(1)'$, implying a degenerate, massive color-octet of coloron (ie. the top-gluon) V_μ^α ($\alpha = 1, 2, \dots, 8$) and a color-singlet heavy Z'_μ . The gluon A_μ^α and coloron V_μ^α (the SM $U(1)_Y$ field B_μ and the $U(1)'$ field Z'_μ), are then defined by orthogonal rotations with mixing angle θ (θ'):

$$\begin{aligned} h_1 \sin \theta &= h_2 \cos \theta = g_3, \quad \cot \theta = \frac{h_1}{h_2}, \\ q_1 \sin \theta' &= q_2 \cos \theta' = g_1, \quad \cot \theta' = \frac{q_1}{q_2}, \end{aligned} \quad (2)$$

where h_1 , h_2 , q_1 , and q_2 are coupling constants of $SU(3)_1$, $SU(3)_2$, $U(1)_{Y1}$ and $U(1)_{Y2}$, respectively. And g_3 (g_1) is the $SU(3)_C$ ($U(1)_Y$) coupling constant at Λ_{TC} . In order to select the correct top-quark direction for condensation one usually demand $\cot \theta \gg 1$ and $\cot \theta' \gg 1$.

Both the coloron V and the Z' must be heavier than 1 TeV, according to the experimental data from Fermilab Tevatron[17, 18]. After integrating out the heavy coloron and Z' , the effective four-fermion interactions have the form [19, 20]

$$\mathcal{L}_{eff} = \frac{4\pi}{M_B^2} \left\{ \left(\kappa + \frac{2\kappa_1}{27} \right) \bar{\psi}_L t_R \bar{t}_R \psi_L + \left(\kappa - \frac{\kappa_1}{27} \right) \bar{\psi}_L b_R \bar{b}_R \psi_L \right\}, \quad (3)$$

where $\kappa = (g_3^2/4\pi) \cot^2 \theta$ and $\kappa_1 = (g_1^2/4\pi) \cot^2 \theta'$, and M_B is the mass of coloron V^α and Z' .

The effective interactions of (3) can be written in terms of two auxiliary scalar doublets ϕ_1 and ϕ_2 . Their couplings to quarks are given by [21]

$$\mathcal{L}_{eff} = \lambda_1 \bar{\psi}_L \phi_1 \bar{t}_R + \lambda_2 \bar{\psi}_L \phi_2 \bar{b}_R, \quad (4)$$

where $\lambda_1^2 = 4\pi(\kappa + 2\kappa_1/27)$ and $\lambda_2^2 = 4\pi(\kappa - \kappa_1/27)$. At energies below the Topcolor scale $\Lambda \sim 1$ TeV the auxiliary fields acquire kinetic terms, becoming physical degrees of

freedom. The properly renormalized ϕ_1 and ϕ_2 doublets take the form

$$\phi_1 = \begin{pmatrix} f_{\tilde{\pi}} + \frac{1}{\sqrt{2}}(h_t + i\tilde{\pi}^0) \\ \tilde{\pi}^- \end{pmatrix} \quad (5)$$

and

$$\phi_2 = \begin{pmatrix} \tilde{H}^+ \\ \frac{1}{\sqrt{2}}(\tilde{H}^0 + i\tilde{A}^0) \end{pmatrix} \quad (6)$$

where $\tilde{\pi}^\pm$ and $\tilde{\pi}^0$ are the top-pions, $\tilde{H}^{\pm,0}$ and $\tilde{A}^0$ are the b-pions, h_t is the top-Higgs[20], and $f_{\tilde{\pi}} \approx 50 GeV$ is the top-pion decay constant.

From eq.(4), the couplings of top-pions to t- and b-quark can be written as [8]:

$$\frac{m_t^*}{f_{\tilde{\pi}}} \left[i\bar{t}t\tilde{\pi}^0 + i\bar{t}_R b_L \tilde{\pi}^+ + \frac{m_b^*}{m_t^*} \bar{t}_L b_R \tilde{\pi}^+ + h.c. \right] \quad (7)$$

here, $m_t^* = (1 - \epsilon)m_t$ and $m_b^* \approx 1 GeV$ denote the masses of top and bottom quarks generated by topcolor interactions.

For the mass of top-pions, the current $1 - \sigma$ lower mass bound from the Tevatron data is $m_{\tilde{\pi}} \geq 150 GeV$ [15], while the theoretical expectation is $m_{\tilde{\pi}} \approx (150 - 300 GeV)$ [8]. For the mass of b-pions, the current theoretical estimation is $m_{\tilde{H}^0} \approx m_{\tilde{A}^0} \approx (100 - 350) GeV$ and $m_{\tilde{H}} = m_{\tilde{H}^0}^2 + 2m_t^2$ [17]. For the color-singlet $\pi_1^\pm$ and color-octet $\pi_8^\pm$, the current theoretical estimations are $m_{\pi_1} \geq 50 GeV$ and $m_{\pi_8} \approx 200 GeV$ [22, 4]. In this paper, we conservatively consider a little more wider mass ranges of new scalars:

$$\begin{aligned} m_{\tilde{\pi}} &= (100 \sim 500) GeV, \quad m_{\tilde{H}} = (300 - 1000) GeV, \quad m_{\tilde{H}^0, \tilde{A}^0} = (150 - 600) GeV, \\ m_{\pi_1} &= (50 - 100) GeV, \quad m_{\pi_8} = (100 - 300) GeV. \end{aligned} \quad (8)$$

For $f_{\tilde{\pi}}$ and ϵ , we use $f_{\tilde{\pi}} = (50 - 60) GeV$ and $\epsilon = (0.03 - 0.1)$ [8, 21].

The effective Yukawa couplings of ordinary technipions $\pi_1^\pm$ and $\pi_8^\pm$ to fermion pairs can be found in refs.[4, 22, 23]. The relevant gauge couplings of unit-charged scalars to Z^0 gauge boson are basically model-independent and can be written as [4, 22],

$$Z\pi_i^+ \pi_i^- : \quad -ig \frac{1 - 2 \sin^2 \theta_W}{2 \cos \theta_W} (p^+ - p^-) \cdot \epsilon', \quad (9)$$

$$Z\pi_{8\alpha}^+ \pi_{8\beta}^- : \quad -ig \frac{1 - 2 \sin^2 \theta_W}{2 \cos \theta_W} (p^+ - p^-) \delta_{\alpha\beta} \cdot \epsilon', \quad (10)$$

where θ_W is the Weinberg angle, p^+ and p^- are the momenta of relevant scalars, ϵ' is the polarization vector of Z^0 gauge boson, and π_i denotes the color-singlet scalars $\tilde{\pi}^\pm$, $\tilde{H}^\pm$ and $\pi_1^\pm$, respectively.

2.2 The square-root ansatz and experimental constraints

At low energy, potentially large FCNCs arise when the quark fields are rotated from their weak eigenbasis to their mass eigenbasis, realized by the matrices $U_{L,R}$ for the up-type quarks, and by $D_{L,R}$ for the down-type quarks. When we make the replacements, for example,

$$b_L \rightarrow D_L^{bd} d_L + D_L^{bs} s_L + D_L^{bb} b_L, \quad (11)$$

$$b_R \rightarrow D_R^{bd} d_R + D_R^{bs} s_R + D_R^{bb} b_R, \quad (12)$$

the FCNC interactions will be induced. In TC2-I model, the corresponding flavor changing effective Yukawa couplings are

$$\frac{m_t^*}{f_{\tilde{\pi}}} \left[i\tilde{\pi}^+ (D_L^{bs} \bar{t}_R s_L + D_L^{bd} \bar{t}_R d_L) + i\tilde{H}^+ (D_R^{bs} \bar{t}_L s_R + D_R^{bd} \bar{t}_L d_R) + h.c. \right]. \quad (13)$$

Although there are many discussions about the mixing matrixes in the TC2 models[8, 19, 21, 24], there exist no “standard” mixing matrixes currently. In the literature, authors usually use the “square-root ansatz”: to take the square root of the standard model CKM matrix ($V_{CKM} = U_L^\dagger D_L$) as an indication of the size of realistic mixings. It should be denoted that the square root ansatz must be modified because of the strong constraints from the data of $B^0 - \overline{B}^0$ mixing [21, 24].

In the SM, the B^0 meson mixings are occurred in second order weak interactions. There is evidence for $B^0 - \overline{B}^0$ mixing with $\Delta M_{B_d} = (3.05 \pm 0.12) \times 10^{-10} MeV$ [25], and for $B_s^0 - \overline{B}_s^0$ mixing, with $\Delta M_{B_s} > 6 \times 10^{-9} MeV$ ($CL = 95\%$)[25].

In TC2 models, the neutral scalars $\tilde{H}^0$ and $\tilde{A}^0$ can induce a contribution to the $B_q^0 - \overline{B}_q^0$ ($q = d, s$) mass difference [19, 21]

$$\frac{\Delta M_{B_q}}{M_{B_q}} = \frac{7}{12} \frac{m_t^2}{f_{\tilde{\pi}}^2 m_{\tilde{H}^0}^2} \delta_{bq} B_{B_q} F_{B_q}^2 \quad (14)$$

where M_{B_q} is the mass of B_q meson, F_{B_q} is the B_q -meson decay constant, B_{B_q} is the renormalization group invariant parameter, and $\delta_{bq} \approx |D_L^{bq} D_R^{bq}|$. For B_d meson, using the data of $\Delta M_{B_d} = (3.05 \pm 0.12) \times 10^{-10} MeV$ [25] and setting $f_{\tilde{\pi}} = 50 GeV$, $\sqrt{B_{B_d} F_{B_d}} = 200 MeV$ [1], one has the bound

$$\delta_{bd} \leq 0.76 \times 10^{-7} \quad (15)$$

for $m_{\tilde{H}^0} \leq 600 GeV$. This is an important and strong bound on the product of mixing elements $D_{L,R}^{bd}$. As pointed in ref.[19], if one naively uses the square-root ansatz for *both*

D_L and D_R , the bound (15) is violated by about 2 orders of magnitudes. As shown in ref.[19], the “triangular texture” of the mixing matrix may provide a natural suppression of the effect by producing approximately diagonal D_L or D_R matrices. This will give $\delta_{bd} \approx 0$ and avoids the bound.

Numerically, if we use the square-root ansatz for D_L itself and assume that $D_{L,R}^{bd}/V_{td} = a_{L,R}^{td}$, then the bound (15) can be written in a new form

$$|a_R^{td}| \leq 1.7 \times 10^{-3}, \quad (16)$$

for $a_L^{td} = 0.5$ and $m_{\tilde{H}^0} \leq 600 GeV$. It is obviously a very strong constraint on D_R^{bd} .

For B_s meson, the available data is only a lower bound on Δm_{B_s} [25]:

$$\Delta M_{B_s} > 6 \times 10^{-9} MeV = 19.6 \Delta M_{B_d}. \quad (17)$$

But one can get a reliable estimation of ΔM_{B_s} from its relation with ΔM_{B_d} . In the ratio of B_s and B_d mass differences, many common factors cancel, and we have[25]

$$\Delta M_{B_s} = \Delta M_{B_d} \frac{M_{B_s}}{M_{B_d}} \frac{B_{B_s} F_{B_s}^2}{B_{B_d} F_{B_d}^2} \frac{|V_{tb}^* \cdot V_{ts}|^2}{|V_{tb}^* \cdot V_{td}|^2} = 24.9 \Delta M_{B_d}, \quad (18)$$

where we have used $M_{B_d} = 5.279 GeV$, $M_{B_s} = 5.369 GeV$, and $B_{B_s}/B_{B_d} = 1.01 \pm 0.04$ and $F_{B_s}/F_{B_d} = 1.15 \pm 0.05$ from lattice QCD [26]. Using eqs.(14,18) and assuming $D_{L,R}^{bs}/V_{ts} = a_{L,R}^{ts}$, we have

$$|a_R^{ts}| \leq 1.6 \times 10^{-3}, \quad (19)$$

for $a_L^{ts} = 0.5$ and $m_{\tilde{H}^0} \leq 600 GeV$. From eqs.(17) and (18) we believe that the true value of ΔM_{B_s} should be within the range of $(19.6 - 24.9) \Delta M_{B_d}$. The inclusion of uncertainties of the data and input parameters will weaken the above constraints, but can not change them greatly. It is thus reasonable to expect that both $|a_R^{td}|$ and $|a_R^{ts}|$ can not be larger than 0.01 for $a_L^{tj} \approx 1/2$ ($j = d, s$) and $m_{\tilde{H}^0} \leq 600 GeV$. We conservatively use 0.01 as the upper bound on both a_R^{ts} and a_R^{td} afterwards.

In this paper we assume that all elements of D_L and D_R are real because we do not study the CP violation here. We will consider the following two typical cases in numerical calculation:

- Case-A: Assuming $\epsilon = 0.05$ and $f_{\tilde{\pi}} = 50 GeV$, using the square-root ansatz for D_L : $a_L^{ts} = a_L^{td} = 1/2$, but setting $a_R^{ts} = a_R^{td} = 0.01$.

- Case-B: Assuming $\epsilon = 0.05$ and $f_{\tilde{\pi}} = 50\text{GeV}$, and using the square-root ansatz for both D_L and D_R : $a_L^{tj} = 1/2$, $|a_R^{tj}| = 1/2$ ($j = d, s$).

The Case-A is consistent with the constraints from the data of $B_q^0 - \overline{B}_q^0$ ($q = d, s$) mixing. For Case-B, it violates the constraints (16,19). We still consider the Case-B here in order to see what will happen if we use the popular square root ansatz for both D_L and D_R .

In this paper, we fix the following relevant parameters [1, 25] and use them as the standard input(SIP):

$$\begin{aligned} M_W &= 80.41\text{GeV}, \alpha_{em} = 1/129, \sin^2 \theta_W = 0.23, m_c \equiv \overline{m}_c(m_c) = 1.35\text{GeV}, \\ m_t &\equiv \overline{m}_t(m_t) = 170\text{GeV}, \mu_c = 1.3\text{GeV}, \mu_t = 170\text{GeV}, \\ \Lambda_{\overline{MS}}^{(4)} &= 0.325\text{GeV}, \Lambda_{\overline{MS}}^{(5)} = 0.225\text{GeV}, A = 0.84, \lambda = 0.22, \rho = 0, \eta = 0.36 \quad (20) \end{aligned}$$

where the A, λ, ρ and η are Wolfenstein parameters at the leading order. For $\alpha_s(\mu)$ we use the two-loop expression as given in second paper of ref.[1].

3 Rare K-decays and new physics effects

In this paper we use the ‘‘Penguin Box expansion’’ (PBE) approach [27]. One important advantage of this PBE approach is that the rare K-decays in question depend only on one or two basic, universal, process independent functions. At next-to-leading order(NLO), such functions are $X(x_t)$ and X_{NL}^l for decays $K^+ \rightarrow \pi^+ \nu \overline{\nu}$ and $K_L \rightarrow \pi^0 \nu \overline{\nu}$, and $Y(x_t)$ and Y_{NL} for the short-distance part of the decay $K_L \rightarrow \mu^+ \mu^-$. The functions $X(x_t)$ and $Y(x_t)$ describe the dominant m_t -dependent contributions, while the functions X_{NL}^l and Y_{NL} describe the m_t -independent contributions stemming from internal quarks other than the top quark (usually known as the charm part).

3.1 Rare K-decays in the SM

In the SM, the rare K-decays $K^+ \rightarrow \pi^+ \nu \overline{\nu}$, $K_L \rightarrow \pi^0 \nu \overline{\nu}$ and the short-distance part of $K_L \rightarrow \mu^+ \mu^-$ have been studied in great detail and was summarized for instance in a new review paper [1]. At the leading order(LO), the contributions to the rare K-decays from Z^0 -penguin and W-box diagrams are controlled by the functions $C_0(x_i)$ and $B_0(x_i)$ ($i = u, c, t$), which were evaluated long time ago by Inami and Lim[28]. In recent

years, the NLO corrections are calculated systematically by many authors[1]. The great progress in both the theoretical and experimental investigations enable us now to study the new physics effects on the rare K-decays.

At the NLO level, the effective Hamiltonian for $K^+ \rightarrow \pi^+ \nu \bar{\nu}$, $K_L \rightarrow \pi^0 \nu \bar{\nu}$ and $(K_L \rightarrow \mu^+ \mu^-)_{SD}$ can be written as [1],

$$\mathcal{H}_{eff}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = \frac{\alpha_{em} G_F}{2\sqrt{2}\pi \sin^2 \theta_W} \sum_{l=e,\mu,\tau} [\lambda_t X(x_t) + \lambda_c X_{NL}^l] (\bar{s}d)_{V-A} (\bar{\nu}_l \nu_l)_{V-A}, \quad (21)$$

$$\mathcal{H}_{eff}(K_L \rightarrow \pi^0 \nu \bar{\nu}) = \frac{\alpha_{em} G_F}{2\sqrt{2}\pi \sin^2 \theta_W} \lambda_t X(x_t) (\bar{s}d)_{V-A} (\bar{\nu} \nu)_{V-A} + h.c., \quad (22)$$

$$\mathcal{H}_{eff}((K_L \rightarrow \mu^+ \mu^-)_{SD}) = -\frac{\alpha_{em} G_F}{2\sqrt{2}\pi \sin^2 \theta_W} [\lambda_t Y(x_t) + \lambda_c Y_{NL}] (\bar{s}d)_{V-A} (\bar{\mu} \mu)_{V-A} + h.c., \quad (23)$$

where α_{em} is the electromagnetic fine-structure constant, $G_F = 1.16639 \times 10^{-5} GeV^{-2}$ is the Fermi coupling constant, and $\lambda_i = V_{is}^* V_{id}$ and V_{ij} ($i = u, c, t; j = d, s, b$) are the elements of the CKM mixing matrix in the SM. The functions $X(x_t)$, $Y(x_t)$, X_{NL}^l and Y_{NL} are

$$X(x_t) = C_0(x_t) - 4B_0(x_t) + \frac{\alpha_s}{4\pi} X_1(x_t), \quad (24)$$

$$Y(x_t) = C_0(x_t) - B_0(x_t) + \frac{\alpha_s}{4\pi} Y_1(x_t), \quad (25)$$

$$X_{NL}^l = C_{NL} - 4B_{NL}^{1/2}, \quad (26)$$

$$Y_{NL} = C_{NL} - B_{NL}^{-1/2} \quad (27)$$

where the functions $C_0(x_t)$ and $B_0(x_t)$ are leading top quark contributions through the Z^0 -penguin and W-box diagrams respectively, $X_1(x_t)$ and $Y_1(x_t)$ are NLO QCD corrections, C_{NL} is the Z^0 -penguin part in charm sector, and finally the functions $B_{NL}^{1/2}$ and $B_{NL}^{-1/2}$ are the W-box contributions in the charm sector, relevant for the case of final state neutrinos (leptons) with weak isospin $T_3 = 1/2$ ($-1/2$) respectively.

Using the effective Hamiltonians (21, 22) and summing over the three neutrino flavors we have [1]

$$B(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = \kappa_+ \cdot \left[\left(\frac{Im \lambda_t}{\lambda^5} X(x_t) \right)^2 + \left(\frac{Re \lambda_c}{\lambda} P_0(X) + \frac{Re \lambda_t}{\lambda^5} X(x_t) \right)^2 \right], \quad (28)$$

$$B(K_L \rightarrow \pi^0 \nu \bar{\nu}) = \kappa_L \cdot \left(\frac{Im \lambda_t}{\lambda^5} X(x_t) \right)^2, \quad (29)$$

where $\kappa_+ = 4.11 \times 10^{-11}$ and $\kappa_L = 1.80 \times 10^{-10}$ as given in ref.[1].

For the short-distance part of $K_L \rightarrow \mu^+ \mu^-$, we have

$$B(K_L \rightarrow \mu^+ \mu^-)_{SD} = \kappa_\mu \cdot \left[\frac{Re\lambda_c}{\lambda} P_0(Y) + \frac{Re\lambda_t}{\lambda^5} Y(x_t) \right]^2 \quad (30)$$

where $\kappa_\mu = 1.68 \times 10^{-9}$ [1]. The functions $P_0(X)$ and $P_0(Y)$ in eqs.(28,30) describe the contributions from the charm sector and have been defined in ref.[1],

$$P_0(X) = \frac{1}{\lambda^4} \left[\frac{2}{3} X_{NL}^e + \frac{1}{3} X_{NL}^\tau \right] \quad (31)$$

$$P_0(Y) = \frac{Y_{NL}}{\lambda^4}. \quad (32)$$

The explicit expressions for the functions $C_0(x_t)$, $B_0(x_t)$, $X_1(x_t)$, $Y_1(x_t)$, C_{NL} , $B_{NL}^{1/2}$ and $B_{NL}^{-1/2}$ can be found for instance in refs.[1]. For the convenience of the reader, we present these functions in Appendix A.

3.2 New Z^0 -penguin contributions in the TC2-I model

For the rare K-decays under consideration, the new physics will manifest itself by modifying the functions $X(x_t)$ and $Y(x_t)$, as well as the functions X_{NL} and Y_{NL} in effective Hamiltonians (21,22,23).

The new one-loop diagrams can be obtained from the diagrams in the SM by replacing the internal $W^\pm$ lines with the unit-charged scalar lines, as shown in Fig.1. The color-octet $\pi_8^\pm$ does not couple to the $l\nu$ lepton pairs, and therefore does not present in the box diagrams. For the color-singlet scalars, they do couple to $l\nu$ pairs through box diagrams, but the relevant couplings are strongly suppressed by the lightness of m_l . Consequently, we can safely neglect the tiny contributions from those scalars through the box diagrams.

In ref.[4], we evaluated the new one-loop Z^0 -penguin and W-box diagrams for the induced $d\bar{s}Z$ couplings due to the exchange of unit-charged technipions $\pi_1^\pm$ and $\pi_8^\pm$ in the multiscale walking technicolor model[14]. In this paper we use the same method and follow the same procedure to evaluate the one-loop diagrams induced by top-pions $\tilde{\pi}^\pm$ and b-pions $\tilde{H}^\pm$.

We will use dimensional regularization to regulate all the ultraviolet divergences in the virtual loop corrections and adopt the $\overline{MS}$ renormalization scheme. It is easy to show that all ultraviolet divergences are canceled for scalars $\tilde{\pi}^\pm$, $\tilde{H}^\pm$, $\pi_1^\pm$ and $\pi_8^\pm$ respectively, and therefore the total sum is finite.

By analytical evaluations of the Feynman diagrams, we find the effective $d\bar{s}Z$ vertex induced by the charged top-pion exchange,

$$\Gamma_{Z\mu}^I = \frac{1}{16\pi^2} \frac{g^3}{\cos\theta_W} \sum_j \lambda_j \bar{s}_L \gamma_\mu d_L C_0(\xi_j), \quad (33)$$

with

$$C_0(\xi_j) = \frac{a_L^{ts*} a_L^{td} m_\pi^2}{2\sqrt{2}f_\pi^2 G_F M_W^2} \cdot T1(\xi_j), \quad (34)$$

$$T1(\xi_j) = \left[\frac{\xi_j(-1 - 3\xi_j + 2\sin^2\theta_W(1 + \xi_j))}{8(1 - \xi_j)} - \frac{\xi_j^2 \cos^2\theta_W}{2(1 - \xi_j)^2} \ln \xi_j \right] \quad (35)$$

$$(36)$$

where $\lambda_j = V_{js}^* V_{jd}$, $\xi_t = m_t^{*2}/m_\pi^2$ with $m_t^* = (1 - \epsilon)m_t$, $\xi_j = m_j^2/m_\pi^2$ for $j = c, u$.

For the case of unit-charged b-pion $\tilde{H}^\pm$, we have

$$\Gamma_{Z\mu}^{II} = \frac{1}{16\pi^2} \frac{g^3}{\cos\theta_W} \sum_j \lambda_j \bar{s}_R \gamma_\mu d_R C_0(\eta_j), \quad (37)$$

with

$$C_0(\eta_j) = \frac{a_R^{ts*} a_R^{td} m_H^2}{2\sqrt{2}f_\pi^2 G_F M_W^2} \left[\frac{\eta_j(-1 + \eta_j + 2\sin^2\theta_W(1 + \eta_j))}{8(1 - \eta_j)} + \frac{\eta_j^2 \sin^2\theta_W}{2(1 - \eta_j)^2} \ln[\eta_j] \right] \quad (38)$$

where $\eta_t = m_t^{*2}/m_H^2$, $\eta_j = m_j^2/m_H^2$ for $j = c, u$.

For the case of technipions $\pi_1^\pm$ and $\pi_8^\pm$, we have

$$\Gamma_{Z\mu}^{III} = \frac{1}{16\pi^2} \frac{g^3}{\cos\theta_W} \sum_j \lambda_j \bar{s}_L \gamma_\mu d_L C_0(y_j), \quad (39)$$

$$\Gamma_{Z\mu}^{IV} = \frac{1}{16\pi^2} \frac{g^3}{\cos\theta_W} \sum_j \lambda_j \bar{s}_L \gamma_\mu d_L C_0(z_j), \quad (40)$$

with

$$C_0(y_j) = \frac{m_{\pi_1}^2}{3\sqrt{2}F_\pi^2 G_F M_W^2} \cdot T1(y_j), \quad (41)$$

$$C_0(z_j) = \frac{8m_{\pi_8}^2}{3\sqrt{2}F_\pi^2 G_F M_W^2} \cdot T1(z_j), \quad (42)$$

where $y_t = m_{t1}^2/m_{\pi_1}^2$ and $z_t = m_{t1}^2/m_{\pi_8}^2$ with $m_{t1} = \epsilon m_t$, and $y_j = m_j^2/m_{\pi_1}^2$ and $z_j = m_j^2/m_{\pi_8}^2$ for $j = u, c$.

The new C_0 functions in (34,38,41,42) are just the same kind of functions as the Γ_Z in (2.7) of ref.[28] or the $C_0(x_i)$ in (2.18) of the first paper in ref.[1]. Each new C_0 function describes the contribution to the $d\bar{s}Z$ vertex from the corresponding scalar. In the numerical calculations we will include the new contributions to the rare K-decays by simply adding the new C_0 functions with their standard model counterpart $C_0(x_i)$.

In the above calculations, we used the unitary relation $\sum_{j=u,c,t} \lambda_j \cdot \text{constant} = 0$ wherever possible, and neglected the masses for all external quark lines. We also used the functions $(B_0, B_\mu, C_0, C_\mu, C_{\mu\nu})$ whenever needed to make the integrations, and the explicit forms of these complicated functions can be found, for instance, in the Appendix-A of ref.[29].

When the new contributions from charged scalars are included, the functions $X(x_t)$, $Y(x_t)$, $P_0(X)$ and $P_0(Y)$ appeared in eqs.(28,29,30) should be modified as the following,

$$X_{tot} = X(x_t) + X^{New}, \quad (43)$$

$$Y_{tot} = Y(x_t) + X^{New}, \quad (44)$$

$$P_0(X)_{tot} = P_0(X) + P_0^{New}, \quad (45)$$

$$P_0(Y)_{tot} = P_0(Y) + P_0^{New}, \quad (46)$$

with

$$X^{New} = C_0(\xi_t) + C_0(\eta_t) + C_0(y_t) + C_0(z_t), \quad (47)$$

$$P_0^{New} = \frac{1}{\lambda^4} \left[C_{NL}(\tilde{\pi}^\pm) + C_{NL}(\tilde{H}^\pm) + C_{NL}(\pi_1^\pm) + C_{NL}(\pi_8^\pm) \right], \quad (48)$$

where the function X^{New} describes the correction from the dominant top quark part, while the function P_0^{New} corresponds to the charm part. The new charm part is numerically very small: no more than 2% of the total new contribution. For the convenience of the reader, we present the explicit expressions of functions $C_{NL}(\pi_i)$ ($\pi_i = \tilde{\pi}^\pm, \tilde{H}^\pm, \pi_1^\pm, \pi_8^\pm$) in Appendix B.

In the SM, by using the SIP we have

$$\begin{aligned} X(x_t) &= 1.537, & Y(x_t) &= 1.032, \\ P_0(X) &= 0.412, & P_0(Y) &= 0.155. \end{aligned} \quad (49)$$

For the Case-A, by using the SIP (20) and assuming $m_{\tilde{\pi}} = 100\text{GeV}$ and $m_{\tilde{H}} = 300\text{GeV}$, we have

$$X(\xi_t) = 2.258, \quad P_0^{New}(\tilde{\pi}^\pm) = -0.0146, \quad (50)$$

$$X(\eta_t) = -7 \times 10^{-4}, \quad |P_0^{New}(\tilde{H}^\pm)| \leq 10^{-6}, \quad (51)$$

here it is easy to see that $X(\eta_t)$ and $P_0^{New}(\tilde{H}^\pm)$ are clearly much smaller than $X(\xi_t)$ and $P_0^{New}(\tilde{\pi}^\pm)$, we therefore neglect the contribution from the b-pions $\tilde{H}^\pm$ in Case-A.

For the Case-B, by using the SIP (20) and assuming $m_{\tilde{\pi}} = 100GeV$ and $m_{\tilde{H}} = 300GeV$, we have

$$X(\xi_t) = 2.258, \quad P_0^{New}(\tilde{\pi}^\pm) = -0.0146, \quad (52)$$

$$X(\eta_t) = \pm 1.746, \quad P_0^{New}(\tilde{H}^\pm) = \pm 0.146, \quad (53)$$

where the sign of $X(\eta_t)$ and $P_0^{New}(\tilde{H}^\pm)$ will be determined by the sign of $a_R^{ts*}a_R^{td}$. Both $X(\eta_t)$ and $P_0^{New}(\tilde{H}^\pm)$ will be positive (negative) when the product $a_R^{ts*}a_R^{td}$ is negative (positive). In the following, we use the term Case-B1 and Case-B2 to denote the case of assuming $a_R^{ts*}a_R^{td} = 1/4, -1/4$, respectively. For the Case-B1 (Case-B2), the new contribution from $\tilde{\pi}^\pm$ and $\tilde{H}^\pm$ will cancel (enhance) each other.

In TC2 models, the new contribution to the rare K-decays from ordinary technipions is strongly suppressed by a factor of $(\epsilon f_{\tilde{\pi}}/F_\pi)^2 \sim 10^{-3}$ for $F_\pi \approx 123GeV$ and $\epsilon \approx 0.05$, when compared with that from the top-pions. Numerically, less than 5% of the total new contribution is due to $\pi_1^\pm$ and $\pi_8^\pm$. We therefore use the fixed values of $m_{\pi_1} = 50GeV$ and $m_{\pi_8} = 100GeV$ in the following numerical calculations. For heavier technipions, their contributions will become even smaller.

In the multiscale walking technicolor model[4], on the contrary, the enhancements to the rare K-decays due to $\pi_1^\pm$ and $\pi_8^\pm$ can be as large as 2 ~ 3 orders of magnitudes. The major reason is the big difference in how to generate large top quark mass in different models, which in turn result in very different effective Yukawa couplings. In the TC2 models, $\pi_1^\pm$ and $\pi_8^\pm$ couple to the top quark with strength $\epsilon m_t/F_\pi \approx 0.1$, which is much smaller than the coupling in the MWTCM: $m_t/F_Q \approx 4$ for $F_Q \approx 40GeV$ [4]. And finally the technipions $\pi_1^\pm$ and $\pi_8^\pm$ contribute to the rare K-decays very differently in the MWTCM and the TC2 models.

Fig.2a and Fig.2b show the X functions for the Case-A and Case-B, respectively. The short-dashed line is the standard model prediction $X(x_t) = 1.537$. In Fig.2a, the long-dashed curve shows the function $X(\xi_t)$ and the solid curve corresponds to the function X_{tot} . The positive $X(\xi_t)$ greatly enhances the $X(x_t)$ for light $\tilde{\pi}^\pm$. In Fig.2b, the dotted and solid curve show the function X_{tot} function for Case-B1 and Case-B2, respectively.

Within the range of $\epsilon = 0.03 \sim 0.1$, the X functions basically remain unchanged. For $f_{\tilde{\pi}} = 60 \text{ GeV}$, the functions $X(\xi_t)$ and $X(\eta_t)$ will be decreased by a factor of $(5/6)^2$.

In the following two sections we will present the numerical results for the branching ratios $B(K^+ \rightarrow \pi^+ \nu \bar{\nu})$, $B(K_L \rightarrow \pi^0 \nu \bar{\nu})$ and $B(K_L \rightarrow \mu^+ \mu^-)_{SD}$ with the inclusion of new physics effects, compare the theoretical predictions with the data available currently.

4 Rare decays $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 \nu \bar{\nu}$

4.1 The decay $K^+ \rightarrow \pi^+ \nu \bar{\nu}$

The rare decay $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ is theoretically very clean and the long-distance contributions were known to be negligible[1]. When the new contributions from scalars are included, one finds

$$B(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = \kappa_+ \cdot \left[\left(\frac{Im\lambda_t}{\lambda^5} X_{tot} \right)^2 + \left(\frac{Re\lambda_c}{\lambda} P_0(X)_{tot} + \frac{Re\lambda_t}{\lambda^5} X_{tot} \right)^2 \right] \quad (54)$$

where $\kappa_+ = 4.11 \times 10^{-11}[1]$, the functions X_{tot} and $P_0(X)_{tot}$ are given in (43,45).

Using the SIP (20), and assuming $m_{\tilde{\pi}} = (100 - 500) \text{ GeV}$ and $m_{\tilde{H}} = 300 \text{ GeV}$, we have

$$B(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = \begin{cases} 9.39 \times 10^{-11} & \text{in SM} \\ (3.92 - 0.89) \times 10^{-10} & \text{in Case - A} \\ (1.95 - 0.13) \times 10^{-10} & \text{in Case - B1} \\ (6.59 - 2.34) \times 10^{-10} & \text{in Case - B2} \end{cases} \quad (55)$$

On the experimental side, the first event for $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ has been recently observed by the BNL787 collaboration [6], giving

$$B(K^+ \rightarrow \pi^+ \nu \bar{\nu})_{exp} = 4.2_{-3.5}^{+9.7} \times 10^{-10}, \quad (56)$$

in the ball park of the SM expectations. Further data already collected are expected to increase the sensitivity by more than a factor 2, and there are plans to collect data representing a further large increase in sensitivity.

Fig.3a and Fig.3b show the $m_{\tilde{\pi}}$ dependence of $B(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ in Case-A and Case-B, respectively. In Fig.3, the horizontal band between two solid lines corresponds to the data (56), while the short-dashed line is the standard model prediction. The solid curve

in Fig.3a shows the branching ratio $B(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ when the new contributions are included. In Fig.3b, the lower (upper) solid and short-dashed curve are the branching ratios $B(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ in Case-B1 (Case-B2) for $m_{\tilde{H}} = 300, 1000 GeV$, respectively.

From Fig.3b, an upper bound on $m_{\tilde{\pi}}$ can be read out: $m_{\tilde{\pi}} \leq 285 GeV$ for $m_{\tilde{H}} \leq 1000 GeV$ in Case-B1. This upper bound will be weakened by about $50 GeV$ if we consider uncertainties of other parameters.

From Fig.3 one can see that the theoretical predictions for the branching ratio $B(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ in TC2-I model are now in good agreement with the data (56) for all three cases. The uncertainty of the data is still large. Further improvement of the data will be very helpful to constraint the TC2 models from this decay mode.

4.2 The decay $K_L \rightarrow \pi^0 \nu \bar{\nu}$

In the SM, the rare decay $K_L \rightarrow \pi^0 \nu \bar{\nu}$ is completely dominated by short-distance loop effects with the top quark exchanges and there is no theoretical uncertainties due to m_c, μ_c and $\Lambda_{\overline{MS}}$ present in the decay $K^+ \rightarrow \pi^+ \nu \bar{\nu}$. Consequently this decay mode is even cleaner than $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and is very well suited for the probe of new physics if the experimental data can reach the required sensitivity.

When the new contributions from scalars are included, the branching ratio will be

$$B(K_L \rightarrow \pi^0 \nu \bar{\nu}) = \kappa_L \cdot \left(\frac{Im \lambda_t}{\lambda^5} X_{tot} \right)^2. \quad (57)$$

where $\kappa_L = 1.80 \times 10^{-10}$ [1], function X_{tot} is given in (43).

Using the SIP (20) and assuming $m_{\tilde{\pi}} = (100 - 500) GeV$ and $m_{\tilde{H}} = 300 GeV$, we have

$$B(K_L \rightarrow \pi^0 \nu \bar{\nu}) = \begin{cases} 2.74 \times 10^{-11} & \text{in SM} \\ (1.67 - 0.28) \times 10^{-10} & \text{for Case - A} \\ (0.75 - 0.01) \times 10^{-10} & \text{for Case - B1} \\ (2.95 - 0.91) \times 10^{-10} & \text{for Case - B2} \end{cases} \quad (58)$$

On the experimental side, the KTeV group has recently quoted a preliminary result[30]

$$B(K_L \rightarrow \pi^0 \nu \bar{\nu}) \leq 1.8 \times 10^{-6}, \quad 90\% C.L, \quad (59)$$

and the same group aims at reaching in 1999 a single event sensitivity of 3×10^{-9} [31]. The CLOE experiment in *DAΦNE* can also reach the sensitivity of 10^{-9} in the next few years[32]

Fig.4 shows the $m_{\tilde{\pi}}$ dependence of $B(K_L \rightarrow \pi^0 \nu \bar{\nu})$ in all three cases. The short-dashed line is the standard model prediction. The middle solid curve is the branching ratio in Case-A; while the lower (upper) two curves are the branching ratios in Case-B1 (Case-B2) for $m_{\tilde{H}} = 300, 1000 \text{ GeV}$, respectively. The enhancement can be as large as one order of magnitude for Case-B2.

Although the current bound (59) is still about four orders of magnitudes above the theoretical expectation after including the contributions from new scalars, it is possible to measure this gold-plated decay mode with enough sensitivity to probe the effects of new physics in next few years. Sensitivities around 10^{-11} are the goal of three dedicated experiments which have been recently proposed[31, 33, 34]. A recent proposal[33], for example, aims to make a $\sim 15\%$ measurement of $B(K_L \rightarrow \pi^0 \nu \bar{\nu})$.

5 The decay $K_L \rightarrow \mu^+ \mu^-$

The rare decay $K_L \rightarrow \mu^+ \mu^-$ is a potentially important channel to study the weak interaction within the SM, as well as possible effects of new physics. This decay proceeds through two different mechanisms: a dominant long-distance (LD) part from the two-photon intermediate state and a short-distance (SD) part, which in the SM arises from one-loop Z^0 -penguin and W-box diagrams involving gauge bosons. Since the short-distance part is sensitive to the presence of virtual top quark and other new heavy particles predicted by many new physics models², it offers a window into new physics phenomena.

For the decay $K_L \rightarrow \mu^+ \mu^-$, the full branching ratio can be written generally as follows[35]

$$B(K_L \rightarrow \mu^+ \mu^-) = 2\beta B(K_L \rightarrow \gamma\gamma) \left(\frac{\alpha m_\mu}{\pi M_K} \right)^2 (Re[A]^2 + Im[A]^2) \quad (60)$$

$$Re[A] = A_{SD} + A_{LD} \quad (61)$$

where $\beta = \sqrt{1 - 4m_\mu^2/M_K^2}$, $Im[A]$ denote the absorptive contribution arising from a two-photon intermediate state, and finally A_{SD} and A_{LD} represent the short- and long-distance dispersive contribution, respectively.

²In TC2 models, for example, the virtual unit-charged scalars will appear in the Z^0 -penguin diagrams as shown in Fig.1, and thus provide new contributions to the rare K-decays.

In the SM, the short-distance part of $B(K_L \rightarrow \mu^+ \mu^-)$ is [1]

$$B(K_L \rightarrow \mu^+ \mu^-)_{SD} = (1.23 \pm 0.57) \times 10^{-9}, \quad (62)$$

where the error is dominated by the uncertainty of $|V_{cb}|$. Using the measured branching ratio of $B(K_L \rightarrow \gamma\gamma) = (5.92 \pm 0.12) \times 10^{-4}$ [25], one gets

$$B(K_L \rightarrow \mu^+ \mu^-)_{abs} = (7.07 \pm 0.18) \times 10^{-9}, \quad (63)$$

which is very close to the measured rate [7, 25]

$$B(K_L \rightarrow \mu^+ \mu^-) = (7.2 \pm 0.5) \times 10^{-9}. \quad (64)$$

It is easy to see that the rate $B(K_L \rightarrow \mu^+ \mu^-)$ is almost saturated by the absorptive contribution, leaving only a small room for the coherent sum of the long- and short-distance dispersive contribution,

$$B(K_L \rightarrow \mu^+ \mu^-)_{dis}^{(LD+SD)} = (0.1 \pm 0.5) \times 10^{-9}. \quad (65)$$

Therefore, the magnitude of the total real part $Re[A]$ must be relatively small compared with the absorptive part ³. Such a small total dispersive amplitude can be realized either when the A_{SD} and A_{LD} parts are both small or by partial cancellation between these two parts.

In ref.[35], the authors estimated the dispersive two-photon contribution to the decays $P \rightarrow l^+ l^-$ ($P = \pi^0, \eta, K_L$, and $l = e, \mu$) in the framework of the chiral perturbation theory and large- N_C considerations and found that

$$\begin{aligned} Re[A(P \rightarrow l^+ l^-)] &= \frac{1}{4\beta} \ln^2\left[\frac{1-\beta}{1+\beta}\right] + \frac{1}{\beta} Li_2\left[\frac{\beta-1}{\beta+1}\right] + \frac{\pi^2}{12\beta} + 3 \ln\left[\frac{m_l}{M_\rho}\right] + \chi(M_\rho) \\ &= \begin{cases} 3.2_{-1.0}^{+0.8}, & \text{for } \eta \rightarrow \mu^+ \mu^- \\ 2.9_{-1.0}^{+0.8} - A_{SD}, & \text{for } K_L \rightarrow \mu^+ \mu^- \end{cases} \end{aligned} \quad (66)$$

where $M_\rho = 0.77 GeV$ is the mass of ρ meson, $\chi(M_\rho) = 5.5_{-1.0}^{+0.8}$ is the local contribution determined by fitting the measured ratio $B(\eta \rightarrow \mu^+ \mu^-) = (5.8 \pm 0.8) \times 10^{-6}$ [25]. The relative sign between the short- and long-distance dispersive amplitude in eq.(66) is fixed by the known positive sign of g_8 in the large- N_c limit [36]. From the measured branching

³The data constrain only the size of $Re[A]$, and thus leaves an ambiguity for the sign of $Re[A]$.

ratios of $\eta \rightarrow \mu^+\mu^-$ and $K_L \rightarrow \mu^+\mu^-$ decays, constraints on the short-distance part A_{SD} of the decay $K_L \rightarrow \mu^+\mu^-$ were derived [35]:

$$A_{SD} = \begin{cases} 2.2^{+1.1}_{-1.3}, & \text{for } \text{Re}[A] > 0 \\ 3.6 \pm 1.2, & \text{for } \text{Re}[A] < 0 \end{cases} \quad (67)$$

The first bound is in good agreement with the standard model expectation $A_{SD}^{SM} = 1.8 \pm 0.6$ [35], while the second bound shows a discrepancy of about 1.4σ with the SM expectation. However, the errors of above two bounds could be reduced by improving the measurements of the branching ratio $\eta \rightarrow \mu^+\mu^-$ and $K_L \rightarrow \mu^+\mu^-$. From eqs.(66) and (67), one can see that the short- and long-distance dispersive contributions are comparable in size but cancel each other strongly. Above bounds on A_{SD} can be translated into the constraints on $B(K_L \rightarrow \mu^+\mu^-)_{SD}$ directly,

$$B(K_L \rightarrow \mu^+\mu^-)_{SD} = \begin{cases} (0.2 - 2.8) \times 10^{-9}, & (\text{Bound} - 1 : \text{ for } \text{Re}[A] > 0) \\ (1.5 - 6.0) \times 10^{-9}, & (\text{Bound} - 2 : \text{ for } \text{Re}[A] < 0) \end{cases} \quad (68)$$

Obviously, the bounds are still relatively weak because of the sign ambiguity of $\text{Re}[A]$.

In the SM, the branching ratio $B(K_L \rightarrow \mu^+\mu^-)_{SD}$ is known at the NLO level[1], as given in (62). By comparing the above bounds on $B(K_L \rightarrow \mu^+\mu^-)_{SD}$ with the theoretical predictions after including new physics contributions, one may find useful information on TC2 models.

When the new contributions from scalars in TC2-I model are included, we have

$$B(K_L \rightarrow \mu^+\mu^-)_{SD} = \kappa_\mu \cdot \left[\frac{\text{Re}\lambda_c}{\lambda} P_0(Y)_{tot} + \frac{\text{Re}\lambda_t}{\lambda^5} Y_{tot} \right]^2. \quad (69)$$

where $\kappa_\mu = 1.68 \times 10^{-9}$ [1], the functions Y_{tot} and $P_0(Y)_{tot}$ are given in (44,46). Using the SIP (20) and assuming $m_{\tilde{\pi}} = (100 - 500) \text{ GeV}$ and $m_{\tilde{H}} = 600 \text{ GeV}$, one gets

$$B(K_L \rightarrow \mu^+\mu^-)_{SD} = \begin{cases} 1.25 \times 10^{-9} & \text{in SM} \\ (9.18 - 1.13) \times 10^{-9} & \text{for Case - A} \\ (4.57 - 0.03) \times 10^{-9} & \text{for Case - B1} \\ (15.36 - 3.81) \times 10^{-9} & \text{for Case - B2} \end{cases} \quad (70)$$

The enhancement can reach one order of magnitude.

Fig.5a shows the branching ratio $B(K_L \rightarrow \mu^+\mu^-)_{SD}$ versus the mass $m_{\tilde{\pi}}$ for the Case-A and Case-B1. The short-dashed line is the SM prediction, while the horizontal band

here corresponds to the Bound-1. The upper solid curve in Fig.5a shows the branching ratio $B(K_L \rightarrow \mu^+\mu^-)_{SD}$ in Case-A, and a lower bound $m_{\tilde{\pi}} \geq 280\text{GeV}$ can be read out. The lower three curves in Fig.5a are the ratios $B(K_L \rightarrow \mu^+\mu^-)_{SD}$ in Case-B1 for $m_{\tilde{H}} = 300, 600, 1000\text{GeV}$ respectively.

Fig.5b shows the same thing as the Fig.5a, but the horizontal band here stands for the Bound-2. For the Case-A, the theoretical prediction is consistent with the bound, while the bound-2 prefer light top-pions for Case-B1: $m_{\tilde{\pi}} \leq 230\text{GeV}$ for $m_{\tilde{H}} \leq 1000\text{GeV}$.

Fig.6a and Fig.6b show the same thing as Fig.5a and Fig.5b but for the Case-B2 instead. The upper three curves in Figs.(6a) and (6b) are the ratios $B(K_L \rightarrow \mu^+\mu^-)_{SD}$ in Case-B2 for $m_{\tilde{H}} = 300, 600, 1000\text{GeV}$ respectively. The Case-B2 is completely excluded if $\text{Re}[A] > 0$, but is still allowed if $\text{Re}[A] < 0$ and $m_{\tilde{\pi}} \geq 270\text{GeV}$.

For the decay $K_L \rightarrow \mu^+\mu^-$, there is a strong cancellation between the short- and long-distance dispersive parts. For the Case-A, the enhancement to $B(K_L \rightarrow \mu^+\mu^-)_{SD}$ can reach a factor of 8 for light top-pions. For the Case-B1 (Case-B2), the new contributions from top-pions and b-pions will cancel (enhance) each other, and the resultant enhancement can reach a factor of 4 (12) for light top-pions.

6 Conclusion and discussions

In this paper we calculated the new contributions to the rare FCNC K-decays $K^+ \rightarrow \pi^+\nu\bar{\nu}$, $K_L \rightarrow \pi^0\nu\bar{\nu}$ and $K_L \rightarrow \mu^+\mu^-$ from the new Z^0 -penguin and box diagrams induced by the unit-charged top-pions $\tilde{\pi}^\pm$, b-pions $\tilde{H}^\pm$, and technipions $\pi_1^\pm$ and $\pi_8^\pm$ appeared in the TC2 models. We choose the TC2-I model proposed by Hill [8] as the typical TC2 model to do the analytical and numerical calculations. It is beyond the scope of this paper to consider the detailed differences between TC2 models.

From the analytical evaluations of the one-loop Feynman diagrams, we extract out the new functions $C_0(\pi_i)$ and $C_{NL}(\pi_i)$ ($\pi_i = \tilde{\pi}^\pm, \tilde{H}^\pm, \pi_1^\pm, \pi_8^\pm$) which describe the new Z^0 -penguin contributions due to unit-charged scalars, combine the new functions with their standard model counterparts and use them directly in the calculation of branching ratios. The m_t -dependent term X^{New} in (47) dominates over the m_t -independent term P_0^{New} in (48).

The mixing matrixes D_L and D_R also play an important role for the characteristics and magnitudes of the new contributions. But unfortunately, these new mixing matrices

are really the most undetermined part of the TC2 models. Thanks to the accurate experimental measurement of $B^0 - \overline{B}^0$ mixing, we got some strong constraints (16,19) on the relevant mixing factors. The Case-A is allowed by the constraints (16,19), while the Case-B1 and Case-B2 are also considered for the purpose of comparison and illustration.

For the decay $K^+ \rightarrow \pi^+ \nu \overline{\nu}$, the enhancement to the ratio $B(K^+ \rightarrow \pi^+ \nu \overline{\nu})$ can reach a factor of $2 \sim 7$. The theoretical predictions in TC2-I model are generally agree well with the data (56) for all three cases. Of course, the uncertainty of the data is still very large and further improvement of the data will be very helpful to test or constraint the TC2 models from this decay mode.

The decay $K_L \rightarrow \pi^0 \nu \overline{\nu}$ is the cleanest decay mode among three decay modes in question. The enhancement to the branching ratio $B(K_L \rightarrow \pi^0 \nu \overline{\nu})$ due to the top-pions and b-pion can be as large as one order of magnitude. But the central problem for this decay mode is the very low sensitivity of the available data, which is about four orders of magnitudes above the theoretical expectation. Further improvements of the data will be very essential to find the signals of new physics through this decay mode.

For the decay $K_L \rightarrow \mu^+ \mu^-$, the situation becomes more complicated because of the involvement of the long-distance contributions. After including the additional short-distance part from new physics, the theoretical predictions are still consistent with the data for Case-A and Case-B1. The Case-B2, however, is disfavored by the data. The major obstacles in extracting strong constraints on A_{SD} out of the decay $K_L \rightarrow \mu^+ \mu^-$ is the large uncertainty of A_{LD} , the sign ambiguity of $Re[A]$ as well as the strong cancellation between the short- and long-distance dispersive parts. And therefore improvements in theoretical predictions and the experimental data will be very essential for us to test the new physics effects through this decay mode.

In summary, the unit-charged scalars appeared in TC2 models can provide sizable new contributions to the rare K-decays $K^+ \rightarrow \pi^+ \nu \overline{\nu}$, $K_L \rightarrow \pi^0 \nu \overline{\nu}$ and $K_L \rightarrow \mu^+ \mu^-$ through Z^0 -penguin diagrams. The accurate data of $B^0 - \overline{B}^0$ mixing lead to strong constraints on the size of D_R^{bd} and D_R^{bs} if we use the square-root ansatz for the D_L . Some simple but interesting lower or upper bounds on $m_{\tilde{\pi}}$ are obtained by comparing the theoretical predictions with the relevant data. The TC2-I model is, in general, still consistent with the available data of rare K-decays in question. Further improvement of the data and the theoretical predictions will be very helpful to constraint the TC2 models from the rare K-decays.

ACKNOWLEDGMENT

This work is supported in part by the National Natural Science Foundation of China under the Grant No.19575015, and by the Outstanding Young Teacher Foundation of the Education Ministry of China, as well as the funds from Henan Science and Technology Committee.

Appendix A

In this Appendix, we present the explicit expressions for the functions $B_0(x_t)$, $C_0(x_t)$, $X_1(x_t)$, $Y_1(x_t)$. The functions of $C_0(x_t)$ and $B_0(x_t)$ govern the leading top quark contributions through the Z^0 -penguin and W-box diagrams in the SM, while the functions $X_1(x_t)$ and $Y_1(x_t)$ describe the NLO QCD corrections,

$$B_0(x_t) = \frac{1}{4} \left[\frac{x_t}{1-x_t} + \frac{x_t \ln[x_t]}{(x_t-1)^2} \right] \quad (A1)$$

$$C_0(x_t) = \frac{x_t}{8} \left[\frac{x_t-6}{x_t-1} + \frac{3x_t+2}{(x_t-1)^2} \ln[x_t] \right] \quad (A2)$$

$$X_1(x_t) = -\frac{23x_t + 5x_t^2 - 4x_t^3}{3(1-x_t)^2} + \frac{x_t - 11x_t^2 + x_t^3 + x_t^4}{(1-x_t)^3} \ln[x_t] \\ + \frac{8x_t + 4x_t^2 + x_t^3 - x_t^4}{2(1-x_t)^3} \ln^2[x_t] - \frac{4x_t - x_t^3}{(1-x_t)^2} L_2(1-x_t) + 8x_t \frac{\partial X_0(x_t)}{\partial x_t} \ln[x_\mu] \quad (A3)$$

$$Y_1(x_t) = -\frac{4x_t + 16x_t^2 + 4x_t^3}{3(1-x_t)^2} - \frac{4x_t - 10x_t^2 - x_t^3 - x_t^4}{(1-x_t)^3} \ln[x_t] \\ + \frac{2x_t - 4x_t^2 + x_t^3 - x_t^4}{2(1-x_t)^3} \ln^2[x_t] - \frac{2x_t + x_t^3}{(1-x_t)^2} L_2(1-x_t) + 8x_t \frac{\partial Y_0(x_t)}{\partial x_t} \ln[x_\mu] \quad (A4)$$

where $x_t = m_t^2/m_W^2$, $x_\mu = \mu^2/M_W^2$ with $\mu = \mathcal{O}(m_t)$ and

$$L_2(1-x_t) = \int_1^{x_t} dy \frac{\ln[y]}{1-y}. \quad (A5)$$

For the charm sector in the SM, the C_{NL} is the Z^0 -penguin part and the $B_{NL}^{1/2}$ ($B_{NL}^{-1/2}$) is the box contribution, relevant for the case of final state leptons with $T_3 = 1/2$ ($T_3 = -1/2$):

$$C_{NL} = \frac{x(m)}{32} K_c^{24/25} \left[\left(\frac{48}{7} K_+ + \frac{24}{11} K_- - \frac{696}{77} K_{k33} \right) \left(\frac{4\pi}{\alpha_s(\mu)} + \frac{15212}{1875} (1 - K_c^{-1}) \right) \right]$$

$$\begin{aligned}
& + \left(1 - \ln \frac{\mu^2}{m^2}\right) (16K_+ - 8K_-) - \frac{1176244}{13125}K_+ - \frac{2302}{6875}K_- + \frac{3529184}{48125}K_{33} \\
& + K \left(\frac{56248}{4375}K_+ - \frac{81448}{6875}K_- + \frac{4563698}{144375}K_{33} \right) \Big] \quad (A6)
\end{aligned}$$

where

$$K = \frac{\alpha_s(M_W)}{\alpha_s(\mu)}, \quad K_c = \frac{\alpha_s(\mu)}{\alpha_s(m)}, \quad K_+ = K^{6/25}, \quad K_- = K^{-12/25}, \quad K_{33} = K^{-1/25} \quad (A7)$$

and

$$\begin{aligned}
B_{NL}^{1/2} = \frac{x(m)}{4} K_c^{24/25} \Bigg[& 3(1 - K_2) \left(\frac{4\pi}{\alpha_s(\mu)} + \frac{15212}{1875}(1 - K_c^{-1}) \right) \\
& - \ln \frac{\mu^2}{m^2} - \frac{r \ln r}{1 - r} - \frac{305}{12} + \frac{15212}{625}K_2 + \frac{15581}{7500}K K_2 \Bigg] \quad (A8)
\end{aligned}$$

$$\begin{aligned}
B_{NL}^{-1/2} = \frac{x(m)}{4} K_c^{24/25} \Bigg[& 3(1 - K_2) \left(\frac{4\pi}{\alpha_s(\mu)} + \frac{15212}{1875}(1 - K_c^{-1}) \right) \\
& - \ln \frac{\mu^2}{m^2} - \frac{329}{12} + \frac{15212}{625}K_2 + \frac{30581}{7500}K K_2 \Bigg] \quad (A9)
\end{aligned}$$

here $K_2 = K_{33}$, $m = m_c$, $\mu = \mathcal{O}(m_c)$, $x(m) = m_c^2/M_W^2$, $r = m_l^2/m_c^2(\mu)$ and m_l is the lepton mass.

Appendix B

For the dominant top sector in the TC2-I model, the $C_0(\pi_i)$ ($\pi_i = \tilde{\pi}^\pm, \tilde{H}^\pm, \pi_1^\pm, \pi_8^\pm$) functions have been given in (34–38).

For the charm sector in the TC2-I model, the $C_{NL}(\pi_i)$ ($\pi_i = \tilde{\pi}^\pm, \tilde{H}^\pm, \pi_1^\pm, \pi_8^\pm$) functions take the form

$$\begin{aligned}
C_{NL}(\pi_i) = a_i \frac{m_c^2}{m_W^2} K_c^{24/25} \Bigg[& \left(\frac{48}{7}K_+ + \frac{24}{11}K_- - \frac{696}{77}K_{33} \right) \left(\frac{4\pi}{\alpha_s(\mu)} + \frac{15212}{1875}(1 - K_c^{-1}) \right) \\
& + \left(1 - \ln \frac{\mu^2}{m^2}\right) (16K_+ - 8K_-) - \frac{1176244}{13125}K_+ - \frac{2302}{6875}K_- + \frac{3529184}{48125}K_{33} \\
& + K_{\pi_i} \left(\frac{56248}{4375}K_+ - \frac{81448}{6875}K_- + \frac{4563698}{144375}K_{33} \right) \Bigg]. \quad (B1)
\end{aligned}$$

with

$$K_c = \frac{\alpha_s(\mu)}{\alpha_s(m_c)}, \quad K_{\pi_i} = \frac{\alpha_s(m_{\pi_i})}{\alpha_s(\mu)}, \quad K_+ = (K_{\pi_i})^{6/25},$$

$$K_- = (K_{\pi_i})^{-12/25}, \quad K_{33} = (K_{\pi_i})^{-1/25}, \quad \mu = \mathcal{O}(m_c). \quad (B2)$$

For top-pion $\tilde{\pi}^\pm$ and b-pion $\tilde{H}^\pm$ we have

$$a_1 = \frac{a_L^{ts*} a_L^{td}}{64 \sqrt{2} f_{\tilde{\pi}}^2 G_F}, \quad K_{\tilde{\pi}} = \frac{\alpha_s(m_{\tilde{\pi}})}{\alpha_s(\mu)}, \quad a_2 = \frac{a_R^{ts*} a_R^{td}}{8 \sqrt{2} f_{\tilde{\pi}}^2 G_F}, \quad K_{\tilde{H}} = \frac{\alpha_s(m_{\tilde{H}})}{\alpha_s(\mu)}. \quad (B3)$$

For ordinary technipions $\pi_1^\pm$ and $\pi_8^\pm$ we have

$$a_3 = \frac{1}{96 \sqrt{2} F_\pi^2 G_F}, \quad K_{\pi_1^\pm} = \frac{\alpha_s(m_{\pi_1})}{\alpha_s(\mu)}, \quad a_4 = \frac{1}{12 \sqrt{2} F_\pi^2 G_F}, \quad K_{\pi_8^\pm} = \frac{\alpha_s(m_{\pi_8})}{\alpha_s(\mu)}. \quad (B4)$$

References

- [1] A. J. Buras and R.Fleischer, hep-ph/9704376, to appear in *Heavy Flavor II*, ed. A.J.Buras and M.Lindner (World Scientific, Singapore); G.Buchalla, A.J.Buras and M.E.Lautenbacher, Rev.Mod.Phys. **68**, 1125(1996).
- [2] M.Misiak, S.Pokorski and J.Rosiek, hep-ph/9703442, to appear in "Heavy Flavor II", ed. A.J.Buras and M.Lindner (World Scientific, Singapore).
- [3] A.J. Buras, A. Remanino and L.Silverstrini, Nucl.Phys. B520, 3(1998); G.Valencia, Nucl.Phys. B517, 33(1998); Gi-Chol Cho, Eur.Phys. J. C5, 525(1998).
- [4] Z. J. Xiao, L.X. Lü, H.K. Guo and G.R. Lu, Eur. Phys. J. C7, 487(1999).
- [5] C.D. Lü and Z.J. Xiao, Phys.Rev. D53, 2529(1996); G.R. Lu, Z.H. Xiong and Y.G. Cao, Nucl. Phys. B487, 355(1997); Z.J. Xiao *et al*, Commun.Theor.Phys. 29(1998)431; Z. J. Xiao, L.X. Lü, H.K. Guo and G.R. Lu, Chin. Phys. Lett. 16(1999)88.
- [6] S. Adler et al., Phys. Rev. Lett. **79**, 2204(1997).
- [7] A.P.Heinson *et al.*, Phys. Rev. **D51**, 985(1995); T. Akagi *et al.*, Phys. Rev.Lett. **67**, 2618(1991); T. Akagi *et al.*, (KEK 137), Phys. Rev. **D51**, 2061(1995).
- [8] C.T. Hill, Phys. Lett. **345B**, 483(1995); K.Lane and E.Eichten, *ibid*, 352, 382(1995).
- [9] S. Weinberg, Phys. Rev. **D13**, 974(1976); **19**, 1277(1979); L.Susskind, *ibid*, **20**, 2619(1979).
- [10] E. Farhi and L.Susskind, Phys. Rev. **D20**, 3404(1979).
- [11] S.Dimopoulos and L. Susskind, Nucl. Phys. **B155**, 237(1979); E.Eichten and K.Lane, Phys. Lett. **90B**, 125(1980).
- [12] B. Holdom, Phys. Rev. **D24**, 1441(1981); T. Appelquist, D. Karabali, and L.C.R. Wijewardhana, Phys. Rev. Lett. **57**, 957(1986).
- [13] M.Luty and R.Sundrum, Phys. Rev. Lett. **70**, 529(1993); R.Sundrum and S. Hsu, Nucl.Phys.**B391**, 127(1993); T.Appelquist and J.Terning, Phys.Lett. **315B**, 139(1993).

- [14] K.Lane and E.Eichten, Phys.Lett. **222B**, 274(1989); K.Lane and M.V. Ramana, Phys.Rev. **D44**, 2678(1991).
- [15] K.Lane, presented at the 28th International Conference on High Energy Physics, Warsaw (July 1996), ICHEP 96:367-378; G. Burdman, hep-ph/9802232.
- [16] K.Lane, Phys. Lett. B433, 96(1998).
- [17] G. Burdman, D. Kominis, Phys. Lett. **B403**, 101(1997); G. Burdman, Phys. Lett. **B409**, 443(1997).
- [18] Y. Su, F. Bonini and D. Kominis, Phys. Rev. Lett. **79**, 4075(1997).
- [19] G.Buchalla, G. Burdman, C.T.Hill and D. Kominis, Phys.Rev. **D53**, 5185(1996).
- [20] G.Burdman, hep-ph/9611265.
- [21] D. Kominis, Phys. Lett. **B 358**, 312(1995).
- [22] E.Eichten, I.Hinchliffe, K.Lane and C.Quigg, Rev.Mod.Phys. **56**, 759 (1984); Phys. Rev. **D34**, 1547(1986); E.Eichten and K.Lane, Phys.Lett. **327B**, 129(1994).
- [23] J.Ellis, M.K.Gaillard, D.V.Nanopoulos and P.Sikivie, Nucl.Phys. **B182**, 505(1981).
- [24] K.Lane, Phys. Rev. **D54**, 2204(1996).
- [25] Particle Data Group, C. Caso *et al.*, Eur. Phys. J. **C3**, 1(1998).
- [26] J.M.Flynn, plenary talk at the 28th International Conference on High Energy Physics, Warsaw, July 25-31, 1996 and hep-lat/9611016.
- [27] G.Buchalla, A.J.Buras and M.K.Harlander, Nucl.Phys. **B349**, 1(1991).
- [28] T.Inami and C.S. Lim, Prog.Theor.Phys. **65**(1981)297.
- [29] P.Cho and B.Greistein, Nucl.Phys. **B365**, 365(1991).
- [30] R. Ben-David, *XVI International Workshop on Weak Interactions and Neutrinos*, Capri (1997).
- [31] E.Chen *et al.*, FERMILAB-PUB-97-321-E, hep-ex/9709026(1997).

- [32] F.Bossi, G. Colangelo and G. Isidori, Eur.Phys. J. C6, 109(1999).
- [33] I.-H Chiang *et al.*, AGS Proposal 926(1996).
- [34] T.Inagaki *et al.*, KEK-96/13(1996);
- [35] D.Gómez Dumm and A. Pich, Phys.Rev.Lett. 80, 4633(1998).
- [36] A. Pich and E. de Rafael, Phys.Lett. **B374**, 186(1996).

Figure Captions

Fig.1: The new Z^0 -penguin and box diagrams induced by the internal exchanges of the unit-charged scalars $\pi_1^\pm$, $\pi_8^\pm$, $\tilde{\pi}^\pm$ and $\tilde{H}^\pm$. The dashed lines are scalars and the u_j stands for the up-type quarks (u, c, t).

Fig.2: Plots of the X functions versus the mass $m_{\tilde{\pi}}$ in TC2-I model. For more details see the text.

Fig.3: The branching ratio $B(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ in TC2-I model as a function of $m_{\tilde{\pi}}$ for the Case-A and Case-B. The short-dashed line is the SM prediction, while the horizontal band shows the data (56). The solid curve in (3a) is the branching ratio in Case-A, while the lower (upper) two curves in Fig.3b show the ratios in Case-B1 (Case-B2) for $m_{\tilde{H}} = 300, 1000 GeV$, respectively.

Fig.4: The branching ratio $B(K_L \rightarrow \pi^0 \nu \bar{\nu})$ in TC2-I model as functions of $m_{\tilde{\pi}}$. The short-dashed line is the SM prediction. The long-dashed curve is the ratio in Case-A, while the lower (upper) two curves show the ratios in Case-B1 (Case-B2) for $m_{\tilde{H}} = 300, 1000 GeV$, respectively.

Fig.5: The branching ratio $B(K_L \rightarrow \mu^+ \mu^-)_{SD}$ as a function of $m_{\tilde{\pi}}$ for Case-A and Case-B1. The short-dashed line is the SM prediction. The horizontal band in Fig.5a (Fig.5b) corresponds to the Bound-1 (Bound-2). The upper solid curve stands for the ratio in Case-A, while the lower three curves show the ratios in Case-B1 for $m_{\tilde{H}} = 300, 600, 1000 GeV$, respectively.

Fig.6: The same as in Fig.5, but for the Case-B2.




Fig.1

